arXiv:1211.0370v1 [quant-ph] 2 Nov 2012

Experimental test of universal joint measurement uncertainty relations

Morgan M. Weston, Michael J. W. Hall, Matthew S. Palsson, Howard M. Wiseman and Geoff J. Pryde
Centre for Quantum Computation and Communication Technology (Australian Research Council), Centre for Quantum Dynamics, Griffith University, Brisbane, QLD 4111, Australia

The principle of complementarity is fundamental to quantum mechanics, and restricts the degrees of accuracy with which incompatible quantum observables can be jointly measured. Despite popular conception, the well-known Heisenberg uncertainty relation does not quantify this principle. Here we report the first experimental investigation of universally valid complementarity relations — also known as joint measurement uncertainty relations — and also derive a new, improved, complementarity relation. Our tests exploited Einstein-Podolsky-Rosen-type correlations between two photonic qubits, to jointly measure incompatible observables for one of them. We demonstrate not only that the universally valid relations always hold, but that practical situations exist where the product of the measurement inaccuracies is sufficiently low to violate a widely used, but not universally valid, relation.

Niels Bohr repeatedly emphasised that the fundamental distinction between quantum and classical mechanics is the principle of complementarity, which states that the experimental arrangements required to accurately measure two different observables are, in general, incompatible [1]. Indeed, in replying to the famous critique by Einstein, Podolsky and Rosen on the completeness of the quantum theory [2], Bohr stated that *"it is only the mutual exclusion of any two experimental procedures, permitting the unambiguous definition of complementary physical quantities, which provides room for new physical laws"* [3].

It is commonly thought that Bohr's complementarity principle is captured by Heisenberg-type uncertainty relations such as $\Delta Q \Delta P \geq \hbar/2$, where ΔQ and ΔP are, respectively, the predicted spreads of perfect position and momentum measurements, arising from the underlying position and momentum spreads of a particle's wavefunction. However, this relation does not reflect nor quantify the property that the mutual incompatibility of such perfect measurements makes it impossible to carry them out at the same time. By contrast, a *complementarity relation* should restrict the degree of accuracy with which the position and momentum can be measured simultaneously. More generally, it should imply that no joint measurement procedure can generate perfectly accurate estimates of two incompatible observables.

If two observables A and B are not jointly measurable because they do not commute, one may still jointly measure estimates, $A_{\rm est} \approx A$ and $B_{\rm est} \approx B$, where $A_{\rm est}$ and $B_{\rm est}$ *do* commute. The key question then concerns the quality of these jointly-measurable estimates: how close are they to the original observables of interest? A natural measure of inaccuracy, for an estimate $A_{\rm est}$ of some observable A, is the root mean square error $\epsilon(A_{\rm est}) := \langle (A - A_{\rm est})^2 \rangle^{1/2}$ [4, 5]. This measure vanishes for a perfect estimate, and more generally quantifies the degree to which the estimate statistically deviates from the quantity being estimated. It was shown by Arthurs and Kelly, in 1965, that $\epsilon(Q_{\rm est})\,\epsilon(P_{\rm est}) \geq \hbar/2$ for any joint estimates of position and momentum which are *globally unbiased*, i.e., for which the average values of the estimates are equal to the average values of the corresponding observables for every state [6]. Thus, the more accurately a globally unbiased measurement can estimate the position, the less accurately it can estimate the momentum, and vice versa. The Arthurs-Kelly relation was subsequently generalised to the complementarity relation [5, 7]

$$\epsilon(A_{\rm est})\,\epsilon(B_{\rm est}) \geq \frac{1}{2}C, \tag{1}$$

for globally unbiased joint estimates of observables A and B. Here $C := |\langle [A, B] \rangle|$, and is nonzero only when A and B are incompatible. Other complementarity relations — also called joint measurement uncertainty relations — have been given [8–13], but again are only valid for restricted classes of measurements.

The abovementioned complementarity relations not only lack universal validity but, most importantly, they do not even apply in cases of considerable physical interest. For example, Einstein, Podolsky and Rosen (EPR) based their argument for the incompleteness of quantum theory on the case of two particles having very well defined relative positions and total momenta [2]. In this case, one can make very accurate joint estimates of the position and momentum of the first particle via, for example, a position measurement on the first particle and a momentum measurement on the second particle. Thus, the Arthurs-Kelly relation (1) clearly fails in the EPR scenario.

Indeed, it is only relatively recently that universally valid complementarity relations have been given, holding for arbitrary joint estimates of two observables in any measurement scenario. In particular, following related work by Ozawa on measurement–disturbance relations [14], Hall showed that the inaccuracies of any estimates generated by a joint measurement of observables A and B must satisfy [15]

$$\epsilon(A_{\rm est})\,\epsilon(B_{\rm est}) + \epsilon(A_{\rm est})\,\Delta B_{\rm est} + \Delta A_{\rm est}\,\epsilon(B_{\rm est}) \geq \frac{1}{2}C, \tag{2}$$

where $\Delta A_{\rm est}$ and $\Delta B_{\rm est}$ denote the spreads of the estimates. Thus, the inaccuracies cannot both be zero for incompatible observables A and B. This complementarity relation generalises (1) to arbitrary joint estimates, and is saturated in the EPR scenario [15]. Hall also determined the form of the *optimal* estimates of A and B in any given measurement scenario, i.e., those functions of the measurement outcomes that yield the smallest possible inaccuracies $\epsilon(A_{\rm est})$ and $\epsilon(B_{\rm est})$ [15]. Ozawa has given the formally similar relation [16]

$$\epsilon(A_{\rm est})\,\epsilon(B_{\rm est}) + \epsilon(A_{\rm est})\,\Delta B + \Delta A\,\epsilon(B_{\rm est}) \geq \frac{1}{2}C. \tag{3}$$

This differs from (2) in not depending on the spreads of the estimates *per se*, but on the spreads of A and B. The Hall relation is stronger than the Ozawa relation for the case of optimal estimates, since $\Delta A_{\rm est} \leq \Delta A$ for such estimates [15].

Here we provide the first experimental test of universal complementarity relations, in an EPR-type scenario for which the Arthurs-Kelly relation (1) is shown to be violated. Moreover, a new universal complementarity relation, which is stronger than both the Hall and Ozawa relations for optimal estimates, is proved here and similarly verified.

A new universal complementarity relation.

In addition to experimentally investigating complementarity relations (1)–(3), we also tested the new relation

$$\epsilon(A_{\rm est})\,\frac{\Delta B_{\rm est} + \Delta B}{2} + \epsilon(B_{\rm est})\,\frac{\Delta A_{\rm est} + \Delta A}{2} \geq \frac{1}{2}C, \tag{4}$$

which we have derived and which is proved in appendix A. Note that, unlike (1)–(3), it does not include an $\epsilon(A_{\rm est})\epsilon(B_{\rm est})$ term. Nevertheless, it similarly implies that $\epsilon(A_{\rm est})$ and $\epsilon(B_{\rm est})$ cannot both be zero for incompatible A and B.

Relation (4) is typically stronger than both the Hall and Ozawa relations, and is always stronger in the case that optimal estimates of A and B are made, in any given measurement scenario, as shown in Appendix A.

Measuring the inaccuracies.

To test the validity of complementarity relations (1)–(4), it is necessary to experimentally determine the inaccuracies $\epsilon(A_{\rm est})$ and $\epsilon(B_{\rm est})$. This problem has been previously discussed in context of measurement–disturbance uncertainty relations, which also involve the inaccuracy of an estimate. Ozawa proposed an approach in which $\epsilon(A_{\rm est})$ is determined from the statistics of (separate) measurements of A and $A_{\rm est}$, on each of three different states of the system (defined by A and the state of interest) [17]. Lund and Wiseman proposed an approach not requiring preparation of additional states, in which a weak measurement of A is made prior to the measurement of $A_{\rm est}$ [18], allowing $\epsilon(A_{\rm est})$ to be determined from the corresponding weak-valued joint probability distribution. These two proposals have been recently applied to the verification of measurement–disturbance uncertainty relations in Refs. [19] and [20] respectively.

Our approach is a generalisation of the Lund-Wiseman proposal, in which the weak measurement interaction is replaced by one of arbitrary strength. To introduce the basic concept, consider two quantum observables K and L having respective eigenvalue decompositions $K = \sum_k \kappa_k K_k$ and $L = \sum_l \lambda_l L_l$. The corresponding 'Margenau-Hill' joint quasiprobability distribution is defined by $p_{\rm MH}(k,l) := \langle K_k L_l + L_l K_k\rangle/2$ [21]. It can take negative values for non-commuting observables, but is normalised and has marginal distributions corresponding to the probability distributions of K and L. The mean square deviation between K and L can be calculated [18, 22] from the Margenau-Hill distribution via

$$\langle (K-L)^2\rangle = \sum_{k,l} (\kappa_k - \lambda_l)^2\, p_{\rm MH}(k,l). \tag{5}$$

Further, if K is an observable prior to the application of an arbitrary quantum evolution (including measurement), and L an observable subsequent to its application, then the Margenau-Hill distribution itself can be determined experimentally via an initial weak measurement of the individual K_k projectors postselected on a final measurement of L [18]. Recalling that $\epsilon(A_{\rm est}) = \langle (A - A_{\rm est})^2\rangle^{1/2}$, this allows the inaccuracies of joint estimates of A and B to be experimentally determined from weak measurements of the eigenprojectors of A and B (in separate experiments), postselected on the joint measurement. This approach also applies if L is replaced by any positive-operator-valued measure $\{L_l\}$ having assigned outcome values $\{\lambda_l\}$ [18].

Our generalization of the Lund-Wiseman method relaxes the requirement that the initial measurement interaction be weak. Instead, it is only required that the necessary Margenau-Hill distributions can be obtained from the combined statistics of the initial and final measurements. This has the advantage of being simpler to implement experimentally, although at the cost of increasing the inaccuracy of at least one of the joint estimates. Further, for our particular experiment, only one initial measurement rather than two weak measurements is needed to obtain $\epsilon(A_{\rm est})$ and $\epsilon(B_{\rm est})$. The modification is inspired by the contextual-value formalism of Dressel *et al.*, which generalises many properties of weak measurements to arbitrary interaction strengths [23, 24].

In particular, suppose that in place of the initial weak measurement of the projector K_k, the system undergoes an initial measurement $\mathcal{X}$ describable by a set of measurement operators $\{M_m\}$ [25], such that each K_k can be written in the form $K_k = \sum_m \alpha_{km} M_m^\dagger M_m$ (for more general measurements, see Appendix B). The coefficients α_{km} are called the contextual values of K_k for the measurement context defined by $\mathcal{X}$ [23]. It follows that the joint probability distribution for outcome m from measurement $\mathcal{X}$ and l from the final measurement of L is

$$p(m,l) = \langle M_m^\dagger L_l M_m\rangle, \tag{6}$$

and that the Margenau-Hill distribution for K and L is

$$p_{\rm MH}(k,l) = \frac{1}{2}\sum_m \alpha_{km}\langle M_m^\dagger M_m L_l + L_l M_m^\dagger M_m\rangle. \quad (7)$$

We have found that in many cases one can choose $\mathcal{X}$ such that the Margenau-Hill distribution (7) can be directly obtained either from the measured joint distribution (6) or from (an assumed) theoretical description of L. For example, if $[L_l, M_m] = 0$ then $p_{\rm MH}(k,l) = \sum_m \alpha_{km} p(m,l)$. In all such cases the corresponding mean square deviation $\langle\, (K-L)^2\,\rangle$ can be evaluated via Eq. (5).

Experimental setup.

To provide a discrete analogy of the EPR scenario discussed in the introduction, we generated pairs of optical polarisation qubits having high fidelity with the entangled state

$$|\psi\rangle = \cos\gamma|HV\rangle - \sin\gamma|VH\rangle. \quad (8)$$

Here H and V refer to horizontal and vertical polarisation, which are assigned as eigenstates of the Z operator [25]. For $\gamma = \pi/4$ this state is maximally entangled, allowing perfectly accurate joint estimates of the X (diagonal/antidiagonal) and Y (right/left-circular) polarisations for the first qubit (via a measurement of Y on the first qubit and X on the perfectly correlated second qubit), analogous to the EPR example. However, this choice is not suitable for testing the complementarity relations (1)–(3), since $C = 2|\langle Z\rangle| = 2|\cos 2\gamma|$ and so trivially vanishes for $\gamma = \pi/4$. Conversely, the choice $\gamma = 0$ (or $\pi/2$) maximises C, but corresponds to a factorisable state from which no information about the X polarisation of the first qubit can be gained via a measurement on the second qubit. Hence an intermediate choice, $\gamma = \pi/8$, was made for our experiment. Quantum state tomography [26] on the state generated by a spontaneous parametric downconversion source gave a fidelity of 97.4±0.3% with the desired entangled state (see Appendix C).

The X and Y polarisations of the first qubit were jointly estimated using the outcomes of a direct measurement of Y on the first qubit and a measurement of some polarisation observable W of the second qubit (Fig. 1). To allow the corresponding inaccuracies to be determined by the method described above, the first qubit was also subjected to an initial semiweak measurement process $\mathcal{X}$ [27], implemented using the polarisation dependence of Fresnel reflection from a glass slide [24] oriented at an angle $\beta = 75°$.

Horizontal and vertical polarisations were reflected with probabilities r_H and r_V and transmitted with probabilities $t_H = 1 - r_H$ and $t_V = 1 - r_V$ (Fig. 1). Wave plates before and after the slide were adjusted to rotate the basis of the semiweak measurement from Z to X, implying that the corresponding measurement operators M_r and M_t are Hermitian with $M_r^2 = r_H X_+ + r_V X_- = \mathbb{1} - M_t^2$, where $X_\pm$ denotes $\frac{1}{2}(\mathbb{1} \pm X)$ and $\mathbb{1}$ is the unit operator. Values of $r_H = 0.1244 \pm 0.0001$ and $r_V = 0.4645 \pm 0.0011$ were found by direct measurement.



FIG. 1: Concept and implementation of the experiment. A source generates entangled pairs of optical qubits, and a joint measurement is used to simultaneously estimate the X and Y polarisations of the first qubit. The joint measurement comprises a semiweak measurement process $\mathcal{X}$ followed by a Y-polarisation measurement on the first qubit, and a W-polarisation measurement on the second qubit. The outcome of $\mathcal{X}$ is used to determine the inaccuracy, $\epsilon(X_{\rm est})$, of any estimate $X_{\rm est} = f(W)$. The combination of the $\mathcal{X}$ and Y measurements denoted by $\mathcal{Y}$, is used to make a noisy estimate $Y_{\rm est} = \mathcal{Y}$ of the Y polarisation, with readily calculable inaccuracy $\epsilon(Y_{\rm est})$. Note this inaccuracy vanishes in the limit that $\mathcal{X}$ is a perfectly weak measurement of X, with $\mathcal{Y} \equiv Y$ for this case. The process $\mathcal{X}$ uses a glass slide as a polarisation-dependent beamsplitter to make an semiweak measurement of X, and a flip mirror (FM) selects between the reflected and transmitted beams to enable the subsequent measurement of Y. Optical elements are coded by colour in the Figure, with representative elements labelled appropriately. QWP and HWP denote a quarter wave and half wave plate, respectively, FPC a fibre polarisation controller, and GT a Glan-Taylor prism.

The contextual values — which represent the idea of eigenvalues generalised to arbitrary measurements [23] — are labelled α_{xm}, for $x = \pm 1$. These follow as $\alpha_{xm} = \frac{1}{2}[1 \pm x(m_H + m_V)/(m_H - m_V)]$ where $+$ $(-)$ applies to the case $m = t$ $(m = r)$ (see previous section). Since the measurement process only acts on the first qubit, then $[M_m, W_w] = 0$ and the inaccuracy of any estimate of the form $X_{\rm est} = f(W)$ follows via Eqs. (5)–(7) as

$$\epsilon(X_{\rm est})^2 = \sum_{x,m,y,w} [x - f(w)]^2\, \alpha_{xm}\, p(m,y,w), \quad (9)$$

where $p(m, y, w)$ is the joint probability of finding outcomes m, y and w for the binary measurements $\mathcal{X}$, Y and W respectively. Hence, the inaccuracy can be directly calculated from the measured reflectivities and joint probability distribution.

The inaccuracy of the estimate $Y_{\rm est}$ can be inferred by taking into account the effect of $\mathcal{X}$ on the first qubit under the assumption that the Y polarisation measurement depicted in Fig. 1 is accurate. In particular, applying $\mathcal{X}$ before the measurement of Y is equivalent to measuring the positive operator valued measure $\mathcal{Y} = \{\tilde{Y}_y\}$ on the initial state, with $\tilde{Y}_y = \sum_m M_m^\dagger Y_y M_m$ [25]. This simplifies to $\tilde{Y}_y = \kappa/2 + (1-\kappa)Y_y$, where $\kappa := 1 - \sqrt{r_H r_V} - \sqrt{t_H t_V}$. Thus the effect of $\mathcal{X}$ is to add white noise of magnitude κ to the statistics of Y for the original state. Note that the value of κ is a measure of the strength of $\mathcal{X}$, with $\kappa = 0$ corresponding to the 'weak limit' $r_H = r_V$, and $\kappa = 1$ corresponding to a projective measurement of X. The measured values of r_H and r_V above yield $\kappa = 0.0749 \pm 0.008$, moderately above the weak limit $\kappa \to 0$. The Margenau-Hill distribution for Y and $\tilde{Y}$ follows as $p_{MH}(y, \tilde{y}) = \frac{1}{2}\langle Y_y \tilde{Y}_{\tilde{y}} + \tilde{Y}_{\tilde{y}} Y_y \rangle$. The corresponding inaccuracy of the estimate $Y_{\rm est} = \tilde{Y}$ can immediately be calculated via Eq. (5) as

$$\epsilon(Y_{\rm est})^2 = \sum_{y,\tilde{y}} [y - \tilde{y}]^2 \, p_{MH}(y, \tilde{y}) = 2\kappa, \tag{10}$$

irrespective of the input state. Note the inaccuracy vanishes in the weak limit $\kappa \to 0$, as expected.

Experimental results.

The joint probability distribution $p(m, y, w)$ was measured for several choices of the polarisation observable W, corresponding to the Bloch sphere angles $\theta = 90^\circ$ and $\phi = 135^\circ, 157.5^\circ, 180^\circ, 202.5^\circ$, and 225°. Note that $\theta = 90^\circ$ and $\phi = 180^\circ$ corresponds to the observable of the second qubit that is most strongly correlated with the X polarisation of the first qubit, for state $|\psi\rangle$ in (8), in the sense of maximising $\langle X \otimes W \rangle$.

Two types of estimates for X, based on the outcome $w = \pm 1$ of the W measurement, were considered. The first type corresponded to simply estimating X to be w, i.e., $X_{\rm est}^{\rm simple} = W$. The second type of estimate exploited the tomographically-determined state of the source, ρ, to make the optimal estimate of the form $X_{\rm est}^{\rm opt} = f_{\rm opt}(W)$ with [15]

$$f_{\rm opt}(w) := \left\langle X \otimes \frac{\mathbb{1} + wW}{2} \right\rangle_\rho \div \left\langle \mathbb{1} \otimes \frac{\mathbb{1} + wW}{2} \right\rangle_\rho .$$

For example, for $W = -X$, i.e., $\theta = 90^\circ$ and $\phi = 180^\circ$, this gave $f_{\rm opt}(1) = 0.630$ and $f_{\rm opt}(-1) = -0.643$. The corresponding inaccuracies were determined via Eq. (9), and are plotted in Fig. 2. It was also verified that the inaccuracy and spread of $X_{\rm est}^{\rm opt}$ satisfied the Hall inaccuracy–dispersion relation [15]

$$\epsilon(X_{\rm est}^{\rm opt})^2 + (\Delta X_{\rm est}^{\rm opt})^2 = (\Delta X)^2 \tag{11}$$

for optimal estimates (Fig. 2).



FIG. 2: Inaccuracies of simple and optimal estimates of X from a measurement of W, and verification of relation (11) for the optimal estimate. Theoretical curves are given for the inaccuracies $\epsilon(X_{\rm est}^{\rm simple})$ and $\epsilon(X_{\rm est}^{\rm opt})$ (upper and lower dotted lines), and for the spread $\Delta X_{\rm est}^{\rm opt}$ (dashed line). The solid line represents the tomographically determined value of ΔX, with the closely adjacent data points corresponding to the square root of the left hand side of Eq. (11). Error bars not shown are smaller than the size of the markers.

The inaccuracy of the estimate $Y_{\rm est} = \tilde{Y}$ was determined via Eq. (10) as $\epsilon(Y_{\rm est}) = (2\kappa)^{1/2} = 0.387 \pm 002$. Further, the spreads $\Delta X_{\rm est}$ and $\Delta Y_{\rm est}$ were calculated directly from the measurement outcomes, and $\Delta X = 0.998 \pm 0.002$, $\Delta Y = 0.9998 \pm 0.0001$, and $C = 2|\langle Z \rangle| = 1.422 \pm 0.008$ from the tomographically-determined state.

The above data allowed complementarity relations (1)-(4) to be tested, for the two types of estimate of X. The universal complementarity relations (2)-(3) were validated and the Arthurs-Kelly relation violated (Fig. 3). The latter violation was expected for EPR scenarios, as discussed in the introduction. For the case of the optimal estimate of X (lower graph in Fig. 3), the results also verified the new complementarity relation (4) is stronger than the Hall and Ozawa relations (2) and (3), as expected (see Appendix A).

Discussion

We have provided the first experimental verification of universal complementarity relations, that hold for any joint measurement of two observables (Fig. 3). This was done in an EPR-type scenario, for which the Arthurs-Kelly relation (1) is violated. We have, furthermore, derived and verified a new universal complementarity relation, Eq. (4), that is significantly stronger than the previously obtained relations (2) and (3) for the case of optimal estimates, and have also verifed the inaccuracy–dispersion relation (11) for optimal estimates.

It would also be of interest to verify the universal complementarity relations for the case of the quadratures of a single-mode optical fields, for the continuously-valued



FIG. 3: Experimental test of complementarity relations. The upper and lower graphs correspond to the simple and the optimal estimates of X described in the text. Each graph shows, in descending order, the left hand sides of the Hall and Ozawa relations (2) and (3) (indistinguishable in the top graph); the new complementarity relation (4);and the left hand side of the Arthurs-Kelly relation (1). The pink shading indicates the region corresponding to violation of any of these relations. It is seen that the universal complemenarity relations (2)–(4) are satisfied whereas the Arthurs-Kelly relation (1) is strongly violated. Error bars not shown are smaller than the size of the markers. Solid curves are theoretical predictions.

quadrature observables $Q = a + a^\dagger$ and $P(a - a^\dagger)/i$. This would allow the relations to be tested in the context more akin to the original scenario envisaged by Einstein, Podolsky and Rosen [2], and have the advantages that (i) the righthand side of the relations is a fixed constant, independent of the state, in this case; and (ii) the relations can be saturated for Gaussian states [15].

Our work represents an important advance in the quantitative understanding and experimental verification of complementarity, arguably the most important foundational principle of quantum mechanics. This principle underlies many aspects of quantum information technology, ranging from to entanglement verification [28] to quantum dense coding [29] to the security of quantum cryptography [30], and our work could have implications in all these areas.

Acknowledgments

MMW thanks Dylan Saunders, Adam Bennet, Allen Boston and Sacha Kocsis for helpful contributions. MJWH thanks Cyril Branciard for pointing out the difference between the joint uncertainty relations in Refs. [15] and [16], and for stimulating discussions. This research was supported by the ARC Centre of Excellence CE110001027.

Appendix A: Proof of Eq. (4)

Any two joint estimates of observables A and B may be represented by two commuting operators $A_{\rm est}$ and $B_{\rm est}$ on a suitable Hilbert space (via a Naimark extension if necessary) [15, 16]. It follows that $2[A,B] = [A - A_{\rm est}, B + B_{\rm est}] + [A + A_{\rm est}, B - B_{\rm est}]$, and hence, using the triangle inequality, that

$$\begin{aligned} 2|\langle [A,B]\rangle| &= |\langle [A - A_{\rm est}, B] + [A - A_{\rm est}, B_{\rm est}] \\ &\quad + [A, B - B_{\rm est}] + [A_{\rm est}, B - B_{\rm est}]\rangle| \\ &\leq |\langle [A - A_{\rm est}, B]\rangle| + |\langle [A - A_{\rm est}, B_{\rm est}]\rangle| \\ &\quad + |\langle [A, B - B_{\rm est}]\rangle| + |\langle [A_{\rm est}, B - B_{\rm est}]\rangle| . \end{aligned}$$

Applying the Schwarz inequality (which lies behind all of these relations), $|\langle [R,S]\rangle| \leq 2\sqrt{\langle (R-r)^2\rangle \, \langle (S-s)^2\rangle}$, with r and s suitably chosen from 0, $\langle R\rangle$ and $\langle S\rangle$, leads to Eq. (4) as desired. A corresponding measurement-disturbance uncertainty relation will be discussed elsewhere.

Eq. (4) is always stronger than complementarity relations (2) and (3), when optimal estimates $A_{\rm opt}$ and $B_{\rm opt}$ are made from the joint measurement results (corresponding to those functions of the measurement outcomes that yield the smallest possible inaccuracies $\epsilon(A_{\rm est})$ and $\epsilon(B_{\rm est})$). This follows from the property

$$(\Delta A)^2 = (\Delta A_{\rm opt})^2 + \epsilon(A_{\rm opt})^2, \; (\Delta B)^2 = (\Delta B_{\rm opt})^2 + \epsilon(B_{\rm opt})^2 \tag{A1}$$

for optimal estimates [15]. In particular, $\Delta A_{\rm opt} \leq \Delta A$ and $\Delta B_{\rm opt} \leq \Delta B$, and hence Eq. (4) implies Eq. (3) for such estimates. Further, defining $\alpha := \epsilon(A_{\rm opt})/\Delta A$ and $\beta := \epsilon(B_{\rm opt})/\Delta B$, then $0 \leq \alpha, \beta \leq 1$, and the difference of the left hand sides of relations (2) and (4) is $\epsilon(A_{\rm opt})\Delta B \, h(\beta) + \Delta A \, \epsilon(B_{\rm opt}) h(\alpha)$, where $h(x) := (1/2)[\sqrt{1-x^2} - (1-x)]$ is never less than zero. Hence Eq. (4) also implies Eq. (2) for optimal estimates, as claimed.

Appendix B: Inaccuracies and contextual values

In our experiment, the initial measurement process $\mathcal{X}$ in Fig. 1 was describable by a set of measurement operators $\{M_m\}$, so that equations (6) and (7) were applicable to the calculation of the inaccuracies $\epsilon(X_{\rm est})$ and $\epsilon(Y_{\rm est})$. Here we indicate how this method can be extended to general $\mathcal{X}$, via an extension of the definition of contextual values.

In general, $\mathcal{X}$ will be described by some completely positive trace-preserving map $\rho \to \chi(\rho) = \sum_m \chi_m(\rho)$, where a measurement outcome m is obtained with probability $\mathrm{Tr}[\chi_m(\rho)]$ and leaves the system in state $\chi_m(\rho)/\mathrm{tr}[\chi_m(\rho)]$ [31]). Our experiment corresponded to the special case $\chi_m(\rho) = M_m \rho M_m^\dagger$. Suppose that each projection operator K_k for a Hermitian operator K can be written in the form $K_k = \sum_m \alpha_{km} \chi_m^d(\mathbb{1})$, where χ_m^d denotes the dual of χ_m (i.e., $\mathrm{Tr}[A\chi_m(B)] \equiv \mathrm{Tr}[\chi_m^d(A)B]$), and $\mathbb{1}$ is the identity operator. The coefficients α_{km} extend the definition of contextual values in [23], and will again be referred to as the contextual values of K_k for the measurement context defined by $\mathcal{X}$. Eqs. (6) and (7), for the measurement joint probability and Margenau-Hill distributions, then generalise to $p(m,l) = \langle\, \chi_m^d(L_l)\, \rangle$ and $p_{\rm MH}(k,l) = \frac{1}{2}\sum_m \alpha_{km} \langle\, \chi_m^d(\mathbb{1}) L_l + L_l \chi_m^d(\mathbb{1})\, \rangle$, respectively. As before, it is often the case that $\mathcal{X}$ may be chosen such that the Margenau-Hill distribution can be directly obtained either from the measured joint distribution (6) or from (an assumed) theoretical description of L. For example, if $\mathcal{X}$ operates on only one subsystem, and L is an observable of a second subsystem, then χ_m and L may be rewritten in the forms $\chi_m \otimes I$ and $\mathbb{1} \otimes L$, yielding $p_{\rm MH}(k,l) = \sum_m \alpha_{km} p(m,l)$. In any such case the corresponding mean square deviation $\langle\, (K-L)^2\, \rangle$ can be evaluated via Eq. (5), and hence the corresponding inaccuracies determined.

Appendix C: Photon source and error analysis

A 120 mW, linearly polarised continuous wave 404 nm laser diode was used to pump a pair of 0.5 mm thick Bismuth Borate (BiBO) crystals. The crystal pair is oriented such that the optical axes of the pair are crossed in what is known as a 'sandwich', and an additional BiBO crystal is inserted into the pump to precompensate the temporal walkoff between H and V polarisations in the downconversion crystal [32, 33]. When pumped with diagonally polarised light, spontaneous parametric down conversion leads to photon pairs at 808 nm in the maximally entangled $|\Phi^+\rangle$ state. Counting in coincidence with a 3 ns window and 3 nm FWHM interference filters, photon pair rates of ~ 2000 s^{-1} were observed.

Uncertainties predominantly arise from poissonian counting statistics due to the random SPDC generation times, and slow, small thermal drifts in the fibre coupling leading to changes in r_H and r_V during the data collection. The latter effect was characterized by measuring these quantites before and after the data collection, and observing the mean and spread of the values. Uncertainties in directly measured experimental values were combined using standard error propagation formulae to generate uncertainties in values calculated from the data. The small deviations between experiment and theory are attributed to a combination of inexact wave plate settings and other uncompensated small thermal drifts (e.g. fibre birefringence) in the experiment.

[1] Bohr, N. *Atomic Physics and Human Knowledge* (Wiley, New York, 1958), pp. 32-66.

[2] Einstein A., Podolsky, B. & Rosen N., *Phys. Rev.* **47**, 777 (1935).

[3] Bohr, N., *Phys. Rev.* **48**, 696 (1935).

[4] Van Trees, H. L. *Detection, Estimation and Modulation Theory Part 1* (Wiley, New York, 2001), Sec. 2.4.

[5] Ozawa, M., in *Quantum Aspects of Optical Communications*, (eds Bendjaballah, C., Hirota O. & Reynaud, S.) *Lecture Notes in Physics* **378** (Springer, Berlin, 1991), pp. 3-17.

[6] Arthurs E. & Kelly Jr., J. L., *Bell Syst. Tech. J.* **44**, 725 (1965).

[7] Arthurs E. & Goodman, M.S., *Phys. Rev. Lett.* **60**, 2447 (1988).

[8] Wootters, W. K. & Zurek, W. H., *Phys. Rev. D* **19**, 473 (1979).

[9] Martens H. & de Muynck, W., *Found. Phys.* **20**, 357 (1990).

[10] Jaeger, G., Shimony, A. & Vaidman, L., *Phys. Rev. A* **51**, 54 (1995).

[11] Appleby, D. M., *Int. J. Theor. Phys.* **37**, 1491 (1998).

[12] Trifonov, A., Björk, G. & Söderholm, J., *Phys. Rev. Lett.* **86**, 4423 (2001).

[13] Busch, P. & Shilladay C. R., *Phys. Rev. A* **68**, 034102 (2003).

[14] Ozawa, M., *Phys. Rev. A* **67**, 042105 (2003).

[15] Hall, M. J. W., *Phys. Rev. A* **69**, 052113 (2004).

[16] Ozawa, M., *Phys. Lett. A* **320**, 367 (2004).

[17] Ozawa, M., *Ann. Phys.* **311**, 350 (2004).

[18] Lund A. P. & Wiseman, H. M., *New. J Phys.* **12**, 093011 (2010).

[19] Erhart, J., Sponar, S., Sulyok, G., Badurek, G., Ozawa M. & Hasegawa, Y., *Nature Phys.* **8** 185 (2012).

[20] Rozema, L. A., Darabi, L., Mahler, D. H., Hayat, A., Soudagar, Y. & Steinberg, A. M., *Phys. Rev. Lett.* **109**, 100404 (2012).

[21] Margenau, H. & Hill, R. N., *Prog. Theor. Phys.* **26**, 722 (1961).

[22] Wiseman, H. M., *Phys. Lett. A* **311**, 285-291 (2003).

[23] Dressel, J., Agarwal, S. & Jordan, A. N., *Phys. Rev. Lett.* **104**, 240041 (2010).

[24] Dressel, J., Broadbent, C. J., Howell, J. C. & Jordan, A.N., *Phys. Rev. Lett.* **106**, 040402 (2011).

[25] Wiseman, H. M. & Milburn G. J., *Quantum Measurement and Control* (Cambridge University Press, Cambridge, 2010).

[26] White, A. G., Gilchrist, A., Pryde, G. J., O'Brien, J. L., Bremner, M. J., & Langford, N. K., *JOSA B* **24**, 172 (2007).

[27] Pryde, G. J., OBrien, J. L., White A. G., Ralph, T. C. and Wiseman H. M., *Phys. Rev. Lett.* **94**, 220405 (2005).

[28] Buscemi, F., *Phys. Rev. Lett.* **108**, 200401 (2012).

[29] Bennett, C. H. & Wiesner, S. J., *Phys. Rev. Lett.* **69**, 2881 (1992).

[30] Gisin, N., Ribordy, G., Tittel, W. & Zbinden, H., *Rev. Mod. Phys.* **74**, 145 (2002).

[31] Kraus, K., *States, Effects and Operations* (Springer, Berlin, 1983).

[32] Rangarajan, R., Goggin, M. & Kwiat, P., *Opt. Express* **17**, 18920 (2009).

[33] Kwiat, P. G., Waks, E., White, A. G., Appelbaum, I. & Eberhard P. H., *Phys. Rev. A* **60**, R773 (1999).